

RESTRICTED USE PESTICIDE
DUE TO TOXICITY TO FISH AND AQUATIC ORGANISMS
FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS, OR PERSONS UNDER THEIR DIRECT
SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S
CERTIFICATION.



INTRA

Section 18 Emergency Exemption

TO CONTROL RICE DELPHACID IN RICE IN THE STATE OF TEXAS

FOR USE ONLY IN THE FOLLOWING TEXAS COUNTIES:
AUSTIN, BRAZORIA CALHOUN, CHAMBERS, COLORADO, FORT BEND, GALVESTON,
HARRIS, JACKSON, JEFFERSON, LAVACA, LIBERTY, MATAGORDA, VICTORIA,
WALLER, AND WHARTON

QUARANTINE EFFECTIVE DATE: March 3, 2019
EXPIRATION DATE: November 9, 2021

Endigo® ZC
EPA Reg. No. 100-1276

EPA File Number: 19TX03

KEEP OUT OF REACH OF CHILDREN.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY
RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered Endigo ZC label (EPA Reg. No. 100-1276).

This label must be in the possession of the user at the time of application.

Any adverse effects resulting from the use of Endigo ZC under this exemption must be immediately reported to the Texas Department of Agriculture.

CROP USE DIRECTIONS

| Crop | Pest | Rate of Endigo ZC per Acre per Application |
|------|----------------|--|
| Rice | Rice delphacid | 4.5 – 6.0 fl oz |

Use Restrictions:

- To reduce risks to bees due to spray drift, the area within 100 feet of the edge of the field (paddy) must be kept free of flowering weeds.
- **Application Timing:** Apply as required by scouting, usually at intervals of 5 or more days. Base timing and frequency of applications upon insect populations reaching locally determined economic thresholds. Target applications before hopperburn occurs.
- **Maximum Endigo ZC Allowed per Application:** Do not exceed a total of 6.0 fl. oz. of Endigo ZC per acre per application (0.055 lb ai of thiamethoxam and 0.041 lb ai of lambda-cyhalothrin).
- **Maximum Number of Endigo ZC Applications/Maximum Allowed per Year:** Three (highest rate) to four (lowest rate) foliar applications of Endigo ZC may be made by ground or aerial equipment, at a rate of 4.5 – 6.0 fl oz/A/application (0.041 – 0.055 lb thiamethoxam and 0.031 – 0.041 lb lambda-cyhalothrin) not to exceed an annual maximum application rate of 18.5 fl oz/A/year (0.171 lb of thiamethoxam and 0.127 lb of lambda-cyhalothrin). Endigo ZC may be used in the same year thiamethoxam is used as an at-plant seed treatment.
- **Pre-Harvest Interval (PHI):** A 14-day pre-harvest interval, a 5-day retreatment interval (RTI) and a 24-hour restricted entry interval (REI) must be observed.
- **Minimum interval between applications:** 5 days
- **Water Volume:** Apply by ground or air equipment using sufficient water volume to ensure thorough coverage of foliage. **Do not** use less than 2 GPA for aerial applications. **Do not** apply as an ultra-low volume (ULV) spray. Adding an emulsified crop oil (e.g., 1 pt. per acre) when lower aerial application volumes are used is recommended to help improve coverage, reduce evaporation, and improve efficacy.
- **Flood Water:** Do not release flood water within 7 days of an application.
- **Aquaculture:** Do not use treated rice fields for the aquaculture of edible fish and crustacea.

©2019 Syngenta

Endigo® and the Syngenta logo are trademarks of a Syngenta Group Company

Registrant:

Syngenta Crop Protection, LLC

PO Box 18300

Greensboro, NC 27419

Label Code: TX1276172AB0319