

**RESTRICTED USE PESTICIDE:**

Due to **acute eye and dermal toxicity**. For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.

## **Special Local Need – 24(c)**

**EPA SLN Number TX-180002**

**Expires: December 31, 2028**

**FOR USE AND DISTRIBUTION IN THE STATE OF TEXAS.**

### **Ravap E.C. Livestock, Poultry & Premise Insecticide Spray EPA Reg. No. 11556-162**

Keep Out of Reach of Children  
**DANGER**  
**DANGER  POISON**  
**PELIGRO**

Corrosive. Causes irreversible eye damage or skin burns. Harmful if swallowed or absorbed through skin. Do not get in eyes (including animals), on skin, or on clothing.

Premise Use Directions For Livestock Facilities and Equipment

**Use:** Residual Surface Spray

**Insect:** Ticks

**Dilution:** Ravap In Water: 1 gal. in 25 gal. or 5 oz. in 1 gal.

Apply 1 gal. of dilution/500-1000 sq. ft. Thoroughly cover walls, ceilings or other areas where pests rest or congregate in indoor and outside livestock facilities (e.g., barns, sheds, runways, paddocks, stables, traps, pens, feedlots) and equipment (e.g., chutes, head-gates, trailers). Do not spray animals directly with this concentration. Do not allow people or animals to return to treated areas until sprays have thoroughly dried. Do not to apply volume past the point of runoff. Do not spray where runoff to water can occur. Cover or remove all food/feed and water prior to use of this product. Do not apply to hayfields, rangelands, or pastures intended for grazing. Repeat applications at 7 - 10 day intervals.

For use only by US Department of Agriculture, Cattle Fever Tick Eradication Program and Texas Animal Health Commission personnel, or under their direction.

**Elanco US Inc.  
2500 Innovation Way  
Greenfield, IN 46140**